



Phone: 732-390-7750 Fax: 844-683-2244

AsteraCancerCare.org

PATIENT REFERRAL FORM

GENERAL

Patient Name: Last First Middle Pt. DOB: / /

Patient Address:

Patient City: Pt. State: Pt. Zip:

Patient Phone: () - Pt. Height: in. DX: Pt. Weight: lbs.

Patient Allergies:

Insurance: ID#:

Referred by: NPI#:

Office Contact (Required): Office Ph: () -

Office Fax: () -

Office Administrator (Required): Administrator Ph: () -

Astera Infusion Therapy scheduling location request:

- Bridgewater East Brunswick Edison Jersey City Monroe Robbinsville Rutherford Somerset

Required Items/Infusion Process:

Valid/signed prescription including name of medication, exact dosage, and directions (prescription only valid for 6 months, including refills)

- Copy of current insurance card
Recent MD consultation notes: relevant disease being treated must be mentioned in report
Allergies and current medication list
Current labs required for specific medication, as noted on the following page(s) of this form
Has the patient initiated treatment at your office? Yes No

If any future lab tests are needed, please provide patient with a prescription, and have patient bring on day of treatment. Results will be sent to referring physician.

Please note:

- 1. A Letter of Medical Necessity is required for all patients receiving their initial infusion at Astera...
2. Benefit investigations, copay assistance and prior authorizations will be handled by the Astera precert staff if required by the payer. Right to auto-substitute biosimilars based on payer's preference.
3. A pretreatment education session will be provided by an Advanced Practice Provider.
4. Once the infusion is complete, a follow-up notice will be faxed to the referring provider.

Patient Name: _____
Last First Middle

DOB: ____/____/____

Please check the box for medication requested, attach required documentation as noted below, and fax all documents to 844.683.2244. Once all documentation is received, we will contact your patient to schedule appointment. Thanks!

Medication Required Current Lab Results

Note: Progress notes and labs must be completed within the previous 6 months for all new and renewed prescriptions.

- Actemra CBC, Lipid Panel, Liver Function, PPD (prior to initiation)
- Benlysta (IV) None
- Briumvi CBC, Quantitative Serum Immunoglobulin, Prior to initiation – Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
 - Confirm No Vaccinations within 4 Weeks of Therapy
- Cimzia CBC, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
- Cinqair Peak Flow and Other Pulmonary Function Tests
- Cytosan CBC, CMP, UA
- Entyvio Liver Function, PPD (prior to initiation)
- Evenity CMP, Dexa Scan within 2 years Confirm pt. has not had an MI or stroke within previous year
- Fasenra Peak Flow and Other Pulmonary Function Tests
- Ilumya CBC, CMP, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
 - Confirm up to date with vaccines and no live vaccinations within 4 weeks prior to starting therapy or have an active infection. Evaluated for malignancy.
- IVIG Hematocrit, Hemoglobin, IgG Concentrations, Platelets, Renal Function Tests, Urine Output
Provide dose basis in mg/kg. Doses will be rounded to the nearest vial size available.
- Krystexxa G6PD Deficiency, Serum Uric Acid Levels, Confirm Oral Urate Lowering Agent Discontinued
- Leqvio Lipid Panel
- Nucala FEV1, Peak Flow and Other Pulmonary Function Tests
- Nulojix CBC, EBV Serology, Magnesium, Operative Report, Potassium, PPD (prior to initiation)
- Ocrevus CBC, prior to initiation - Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
 - Confirm No Vaccinations within 4 Weeks of Therapy

- Orencia (IV) Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
- Panhematin CMP, Iron Studies, Prior to initiation - Urinary levels of porphobilinogen (PBG), delta aminolevulinic acid (ALA), and total porphyrin
- Prolastin Alpha 1 Proteinase Inhibitor Serum Levels and Lung Function
 - IgA antibodies negative for patient with IgA deficiency
- Radicava None
- Remicade/Inflectra (Biosimilar might be replaced if appropriate)
 - CBC, Liver Function, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
- Rituxan/Riabni/Truxima/Ruxience (CMS approved indications only - Biosimilar might be replaced if appropriate)
 - CBC, prior to initiation - Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
 - Confirm No Vaccinations within 4 Weeks of Therapy
- Saphnelo Up to date with all immunizations before treatment initiation and confirm no live or live attenuated vaccines are given concurrently.
- Simponi Aria (IV) CBC, Liver Function, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
- Skyrizi (IV) Crohn’s Disease Indication only - CBC, CMP (with LFTs), Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)
 - Confirm No Vaccinations within 4 Weeks of Therapy or have an active infection
- Soliris Meningococcal Vaccination
- Stelara (IV) CBC, PPD
- Tezspire FEV1, Peak Flow and Other Pulmonary Function Tests
- Tysabri MRI (MS patients), TOUCH Program Registration
- Vpriv Gene Testing (GBA – Velaglucerase Alfa)
- Vyepti None
- Vyvgart CBC, Anti-AChR Antibody Positive, No Live Vaccines During Therapy
- Xolair Baseline Serum Ige, FEV1, Peak Flow, Other Pulm Function Test (all required for asthma indication only)